

Synthesizing Data: Global Fit (3-Neutrinos)



A. Marrone, Univ. of Bari & INFN

18-24 June 2017 UC Irvine, Irvine, CA, USA

Outline

- Neutrino oscillation parameters:
knowns and unknowns
- Mass ordering from oscillation and
non-oscillation data
- Combined analysis of oscillation
and non-oscillation data
- Conclusions

Outline

- Neutrino oscillation parameters:
knowns and unknowns
- Mass ordering from oscillation and
non-oscillation data
- Combined analysis of oscillation
and non-oscillation data
- Conclusions

Based on

F. Capozzi, E. Di Valentino, E. Lisi, A.M., A. Melchiorri and A. Palazzo Phys.Rev. D95 (2017) no.9, 096014

Precision era in neutrino oscillation phenomenology

Standard 3 ν mass-mixing framework parameters

Precision era in neutrino oscillation phenomenology

Standard 3ν mass-mixing framework parameters

What we known

$$\delta m^2 \quad 2.3\%$$

$$\Delta m^2 \quad 1.6\%$$

$$\sin^2 \theta_{12} \quad 5.8\%$$

$$\sin^2 \theta_{13} \quad 4.0\%$$

$$\sin^2 \theta_{23} \quad \sim 9.6\%$$

Precision era in neutrino oscillation phenomenology

Standard 3 ν mass-mixing framework parameters

What we known

What we still do not know

$$\delta m^2 \quad 2.3\%$$

$$\Delta m^2 \quad 1.6\%$$

$$\sin^2 \theta_{12} \quad 5.8\%$$

$$\sin^2 \theta_{13} \quad 4.0\%$$

$$\sin^2 \theta_{23} \quad \sim 9.6\%$$

Precision era in neutrino oscillation phenomenology

Standard 3 ν mass-mixing framework parameters

What we known

δm^2 2.3%

Δm^2 1.6%

$\sin^2 \theta_{12}$ 5.8%

$\sin^2 \theta_{13}$ 4.0%

$\sin^2 \theta_{23}$ $\sim 9.6\%$


What we still do not know

CP-violating phase δ

Octant of θ_{23}

Mass Ordering $\rightarrow \text{sign}(\Delta m^2)$

$$\Delta m^2 = (\Delta m_{13}^2 + \Delta m_{23}^2)/2$$



Precision era in neutrino oscillation phenomenology

Standard 3ν mass-mixing framework parameters

What we known

δm^2 2.3%

Δm^2 1.6%

$\sin^2 \theta_{12}$ 5.8%

$\sin^2 \theta_{13}$ 4.0%

$\sin^2 \theta_{23}$ $\sim 9.6\%$


What we still do not know

CP-violating phase δ

Octant of θ_{23}

Mass Ordering $\rightarrow \text{sign}(\Delta m^2)$


$$\Delta m^2 = (\Delta m_{13}^2 + \Delta m_{23}^2)/2$$




[We also dont know Dirac/Majorana neutrinos,
Majorana phases, absolute mass scale]

Bounds on single oscillation parameters

Bounds on single oscillation parameters




Bounds on single oscillation parameters




$$N\sigma = \sqrt{\Delta\chi^2} = 1, 2, 3, \dots$$

Bounds on single oscillation parameters



$$N\sigma = \sqrt{\Delta\chi^2} = 1, 2, 3, \dots$$


Bounds on single oscillation parameters



Squared mass differences and mixing angles have both lower & upper bounds at more than 3σ

$$N\sigma = \sqrt{\Delta\chi^2} = 1, 2, 3, \dots$$

Bounds on single oscillation parameters




Squared mass differences and mixing angles have both lower & upper bounds at more than 3σ

Nearly Gaussian uncertainties for
 $(\Delta m^2, \sin^2 \theta_{13})$

and to a lesser extent for
 $(\delta m^2, \sin^2 \theta_{12})$

$$N\sigma = \sqrt{\Delta\chi^2} = 1, 2, 3, \dots$$

Bounds on single oscillation parameters




Squared mass differences and mixing angles have both lower & upper bounds at more than 3σ

Nearly Gaussian uncertainties for
 $(\Delta m^2, \sin^2 \theta_{13})$

and to a lesser extent for
 $(\delta m^2, \sin^2 \theta_{12})$

$$N\sigma = \sqrt{\Delta\chi^2} = 1, 2, 3, \dots$$

Bounds on single oscillation parameters



Squared mass differences and mixing angles have both lower & upper bounds at more than 3σ


Nearly Gaussian uncertainties for
 $(\Delta m^2, \sin^2 \theta_{13})$

and to a lesser extent for
 $(\delta m^2, \sin^2 \theta_{12})$

CP phase: $\delta \sim 1.4\pi$ at best fit
 CP-conserving cases ($\delta = 0, \pi$)
 disfavored at $\sim 2\sigma$ level or more
 Significant fraction of the $[0, \pi]$ range
 disfavored at $> 3\sigma$

$$N\sigma = \sqrt{\Delta\chi^2} = 1, 2, 3, \dots$$

Bounds on single oscillation parameters



Squared mass differences and mixing angles have both lower & upper bounds at more than 3σ

Nearly Gaussian uncertainties for
 $(\Delta m^2, \sin^2 \theta_{13})$


and to a lesser extent for
 $(\delta m^2, \sin^2 \theta_{12})$

CP phase: $\delta \sim 1.4\pi$ at best fit
 CP-conserving cases ($\delta = 0, \pi$)
 disfavored at $\sim 2\sigma$ level or more
 Significant fraction of the $[0, \pi]$ range
 disfavored at $> 3\sigma$

θ_{23} maximal mixing disfavored at
 about more than 2σ level
 best-fit octant flips with mass
 ordering

$$N\sigma = \sqrt{\Delta\chi^2} = 1, 2, 3, \dots$$

Bounds on single oscillation parameters



Squared mass differences and mixing angles have both lower & upper bounds at more than 3σ

Nearly Gaussian uncertainties for
 $(\Delta m^2, \sin^2 \theta_{13})$

and to a lesser extent for
 $(\delta m^2, \sin^2 \theta_{12})$

CP phase: $\delta \sim 1.4\pi$ at best fit
 CP-conserving cases ($\delta = 0, \pi$)
 disfavored at $\sim 2\sigma$ level or more
 Significant fraction of the $[0, \pi]$ range
 disfavored at $> 3\sigma$

θ_{23} maximal mixing disfavored at
 about more than 2σ level
 best-fit octant flips with mass
 ordering

$$N\sigma = \sqrt{\Delta\chi^2} = 1, 2, 3, \dots$$

$$\Delta\chi^2_{\text{IO-NO}} = 3.6$$

To study parameter correlations we combine different data sets:

To study parameter correlations we combine different data sets:

(1) LBL acc + Solar + KamLAND

To study parameter correlations we combine different data sets:

(1) LBL acc + Solar + KamLAND

Solar + KL data provide a measure of $(\delta m^2, \theta_{12})$, and also independent -although weak- constraints on θ_{13} (first "hint" ever...). LBL provide a measure of $(\Delta m^2, \theta_{23})$. The combination gives, by itself, a measure of θ_{13}

To study parameter correlations we combine different data sets:

(1) LBL acc + Solar + KamLAND

Solar + KL data provide a measure of $(\delta m^2, \theta_{12})$, and also independent -although weak- constraints on θ_{13} (first "hint" ever...). LBL provide a measure of $(\Delta m^2, \theta_{23})$. The combination gives, by itself, a measure of θ_{13}

(2) LBL acc + Solar + KamLAND + SBL Reactors

To study parameter correlations we combine different data sets:

(1) LBL acc + Solar + KamLAND

Solar + KL data provide a measure of $(\delta m^2, \theta_{12})$, and also independent -although weak- constraints on θ_{13} (first "hint" ever...). LBL provide a measure of $(\Delta m^2, \theta_{23})$. The combination gives, by itself, a measure of θ_{13}

(2) LBL acc + Solar + KamLAND + SBL Reactors

SBL reactors not only provide the most accurate determination of θ_{13} but also an independent (and increasingly competitive) determination of Δm^2 , not limited by E-reconstruction systematics as in LBL accel.

To study parameter correlations we combine different data sets:

(1) LBL acc + Solar + KamLAND

Solar + KL data provide a measure of $(\delta m^2, \theta_{12})$, and also independent -although weak- constraints on θ_{13} (first "hint" ever...). LBL provide a measure of $(\Delta m^2, \theta_{23})$. The combination gives, by itself, a measure of θ_{13}

(2) LBL acc + Solar + KamLAND + SBL Reactors

SBL reactors not only provide the most accurate determination of θ_{13} but also an independent (and increasingly competitive) determination of Δm^2 , not limited by E-reconstruction systematics as in LBL accel.

(3) LBL acc + Solar + KamLAND + SBL Reactors + Atmospheric

To study parameter correlations we combine different data sets:

(1) LBL acc + Solar + KamLAND

Solar + KL data provide a measure of $(\delta m^2, \theta_{12})$, and also independent -although weak- constraints on θ_{13} (first "hint" ever...). LBL provide a measure of $(\Delta m^2, \theta_{23})$. The combination gives, by itself, a measure of θ_{13}

(2) LBL acc + Solar + KamLAND + SBL Reactors


SBL reactors not only provide the most accurate determination of θ_{13} but also an independent (and increasingly competitive) determination of Δm^2 , not limited by E-reconstruction systematics as in LBL accel.

(3) LBL acc + Solar + KamLAND + SBL Reactors + Atmospheric


Herein: Atmospheric = SK + DeepCore

Knowns: $(\sin^2 \theta_{13}, \Delta m^2)$

Knowns: $(\sin^2 \theta_{13}, \Delta m^2)$




Knowns: $(\sin^2 \theta_{13}, \Delta m^2)$



Progressive reduction of the allowed
 Δm^2 range by adding new data sets

Knowns: $(\sin^2 \theta_{13}, \Delta m^2)$



Progressive reduction of the allowed
 Δm^2 range by adding new data sets

$\sin^2 \theta_{13}$ best measured by reactors

Unknowns: δ , the octant of θ_{23} and Mass Ordering

Unknowns: δ , the octant of θ_{23} and Mass Ordering



δ

θ_{23}

octant

$\Delta\chi^2_{IO-NO}$





Unknowns: δ , the octant of θ_{23} and Mass Ordering



$\Delta\chi^2_{IO-NO}$

+1.1

Unknowns: δ , the octant of θ_{23} and Mass Ordering









$\Delta\chi^2_{IO-NO}$

+1.1

+1.1

Unknowns: δ , the octant of θ_{23} and Mass Ordering



$\Delta\chi^2_{IO-NO}$


+1.1

+1.1


+3.6

Unknowns: CP phase δ

Unknowns: CP phase δ




Unknowns: CP phase δ



θ_{23} octant degeneracy effect on "wavy bands" in the (δ, θ_{13}) plane


Unknowns: CP phase δ



θ_{23} octant degeneracy effect on "wavy bands" in the (δ, θ_{13}) plane

Largely reduced by adding SBL reactor data (which select a preferred octant for given mass ordering). Note also synergy, not tension, between (LBL acc + solar + KamLAND) and (SBL Reactors) data

Unknowns: CP phase δ




θ_{23} octant degeneracy effect on “wavy bands” in the (δ, θ_{13}) plane

Largely reduced by adding SBL reactor data (which select a preferred octant for given mass ordering). Note also synergy, not tension, between (LBL acc + solar + KamLAND) and (SBL Reactors) data


Results in the (δ, θ_{13}) plane corroborated by atmospheric data

Unknowns: θ_{23} octant

Unknowns: θ_{23} octant




Unknowns: θ_{23} octant



In LBL accelerator data, θ_{23} and θ_{13} anticorrelated via appearance data, with the θ_{23} octants largely degenerate via disappearance data. Degeneracy slightly broken by solar + KL data constraints on θ_{13}


Unknowns: θ_{23} octant



In LBL accelerator data, θ_{23} and θ_{13} anticorrelated via appearance data, with the θ_{23} octants largely degenerate via disappearance data. Degeneracy slightly broken by solar + KL data constraints on θ_{13}

SBL reactors θ_{13} is in agreement with solar best fit

Unknowns: θ_{23} octant



In LBL accelerator data, θ_{23} and θ_{13} anticorrelated via appearance data, with the θ_{23} octants largely degenerate via disappearance data. Degeneracy slightly broken by solar + KL data constraints on θ_{13}

SBL reactors θ_{13} is in agreement with solar best fit

Atmospheric data introduce some differences in the relative likelihood of the two octants in NO and IO

Unknowns: Mass Ordering

Unknowns: Mass Ordering

Info from oscillation experiments

Unknowns: Mass Ordering

Info from oscillation experiments

Unknowns: Mass Ordering

Info from oscillation experiments

- Interference between Δm^2 and $\delta m^2 \rightarrow$ Medium baseline experiments

Unknowns: Mass Ordering

Info from oscillation experiments

- Interference between Δm^2 and $\delta m^2 \rightarrow$ Medium baseline experiments
- Matter effects for accelerator or atmospheric neutrinos

Unknowns: Mass Ordering

Info from oscillation experiments

- Interference between Δm^2 and $\delta m^2 \rightarrow$ Medium baseline experiments
- Matter effects for accelerator or atmospheric neutrinos
- Collective effects due to auto-interactions of Supernova neutrinos

Unknowns: Mass Ordering

Info from oscillation experiments

- Interference between Δm^2 and $\delta m^2 \rightarrow$ Medium baseline experiments
- Matter effects for accelerator or atmospheric neutrinos
- Collective effects due to auto-interactions of Supernova neutrinos

Direct searches

Unknowns: Mass Ordering

Info from oscillation experiments

- Interference between Δm^2 and $\delta m^2 \rightarrow$ Medium baseline experiments
- Matter effects for accelerator or atmospheric neutrinos
- Collective effects due to auto-interactions of Supernova neutrinos

Direct searches

Unknowns: Mass Ordering

Info from oscillation experiments

- Interference between Δm^2 and $\delta m^2 \rightarrow$ Medium baseline experiments
- Matter effects for accelerator or atmospheric neutrinos
- Collective effects due to auto-interactions of Supernova neutrinos

Direct searches

- β decay

Unknowns: Mass Ordering

Info from oscillation experiments

- Interference between Δm^2 and $\delta m^2 \rightarrow$ Medium baseline experiments
- Matter effects for accelerator or atmospheric neutrinos
- Collective effects due to auto-interactions of Supernova neutrinos

Direct searches

- β decay
- $0\nu\beta\beta$ decay

Unknowns: Mass Ordering

Info from oscillation experiments



- Interference between Δm^2 and $\delta m^2 \rightarrow$ Medium baseline experiments
- Matter effects for accelerator or atmospheric neutrinos
- Collective effects due to auto-interactions of Supernova neutrinos

Direct searches

- β decay
- $0\nu\beta\beta$ decay
- Cosmology & Astrophysics


JUNO-like prospective sensitivity to mass ordering (*)

Abscissa scales as $T^{1/2} \rightarrow$ linear behaviour for pure statistical errors



Inclusion of energy-scale uncertainties bends the linear rise, but still allows 3σ discrimination after ~ 6 years of data taking. With the inclusion of flux-shape uncertainties: 3σ sensitivity in ~ 10 years

PINGU sensitivity to mass ordering (similarly for ORCA)



Mass ordering discrimination reduced by 2D spectral shape systematics
due to atmospheric fluxes, cross sections, detector response, ...

Parameters sensitive to absolute masses

Parameters sensitive to absolute masses

- β Decay

$$m_\beta = \sqrt{\sum_i |U_{ei}|^2 m_i^2}$$

Parameters sensitive to absolute masses

- β Decay

$$m_\beta = \sqrt{\sum_i |U_{ei}|^2 m_i^2}$$

- $0\nu\beta\beta$

$$m_{\beta\beta} = \left| \sum_i U_{ei}^2 m_i \right| = |c_{13}^2 c_{12}^2 m_1 + c_{13}^2 s_{12}^2 m_2 e^{i\phi_2} + s_{13}^2 m_3 e^{i\phi_3}|$$

Parameters sensitive to absolute masses

- β Decay

$$m_\beta = \sqrt{\sum_i |U_{ei}|^2 m_i^2}$$

- $0\nu\beta\beta$

$$m_{\beta\beta} = \left| \sum_i U_{ei}^2 m_i \right| = |c_{13}^2 c_{12}^2 m_1 + c_{13}^2 s_{12}^2 m_2 e^{i\phi_2} + s_{13}^2 m_3 e^{i\phi_3}|$$

- Cosmology & Astrophysics

$$\Sigma = m_1 + m_2 + m_3$$

Parameters sensitive to absolute masses

- β Decay

$$m_\beta = \sqrt{\sum_i |U_{ei}|^2 m_i^2}$$

- $0\nu\beta\beta$

$$m_{\beta\beta} = \left| \sum_i U_{ei}^2 m_i \right| = |c_{13}^2 c_{12}^2 m_1 + c_{13}^2 s_{12}^2 m_2 e^{i\phi_2} + s_{13}^2 m_3 e^{i\phi_3}|$$

- Cosmology & Astrophysics

$$\Sigma = m_1 + m_2 + m_3$$

Let's focus on Σ and $m_{\beta\beta}$ (if ν are Majorana)

Parameters sensitive to absolute masses

- β Decay

$$m_\beta = \sqrt{\sum_i |U_{ei}|^2 m_i^2}$$

- $0\nu\beta\beta$

$$m_{\beta\beta} = \left| \sum_i U_{ei}^2 m_i \right| = |c_{13}^2 c_{12}^2 m_1 + c_{13}^2 s_{12}^2 m_2 e^{i\phi_2} + s_{13}^2 m_3 e^{i\phi_3}|$$

- Cosmology & Astrophysics

$$\Sigma = m_1 + m_2 + m_3$$

Let's focus on Σ and $m_{\beta\beta}$ (if ν are Majorana)

- Oscillation data strongly correlate $(\Sigma, m_{\beta\beta})$
- Cosmological data constrain Σ
- $0\nu\beta\beta$ data constrain $m_{\beta\beta}$

Parameters sensitive to absolute masses

- β Decay

$$m_\beta = \sqrt{\sum_i |U_{ei}|^2 m_i^2}$$

- $0\nu\beta\beta$

$$m_{\beta\beta} = \left| \sum_i U_{ei}^2 m_i \right| = |c_{13}^2 c_{12}^2 m_1 + c_{13}^2 s_{12}^2 m_2 e^{i\phi_2} + s_{13}^2 m_3 e^{i\phi_3}|$$

- Cosmology & Astrophysics

$$\Sigma = m_1 + m_2 + m_3$$

Let's focus on Σ and $m_{\beta\beta}$ (if ν are Majorana)

- Oscillation data strongly correlate $(\Sigma, m_{\beta\beta})$
- Cosmological data constrain Σ
- $0\nu\beta\beta$ data constrain $m_{\beta\beta}$



Add up $\chi_{\text{osc}}^2 + \chi_\Sigma^2 + \chi_{\beta\beta}^2$ in the $(\Sigma, m_{\beta\beta})$ plane

When deriving parameter bounds, two possible strategies

When deriving parameter bounds, two possible strategies



- (1) Take NO and IO as two alternative hypotheses

When deriving parameter bounds, two possible strategies



(1) Take NO and IO as two alternative hypotheses



When deriving parameter bounds, two possible strategies





(1) Take NO and IO as two alternative hypotheses

(2) Minimize over any ordering taking into account the offset between the two alternative hypotheses

When deriving parameter bounds, two possible strategies




(1) Take NO and IO as two alternative hypotheses




(2) Minimize over any ordering taking into account the offset between the two alternative hypotheses

Oscillations independent on the absolute mass scale but give rise to a lower bound on Σ to which cosmological data are sensitive




Oscillations independent on the absolute mass scale but give rise to a lower bound on Σ to which cosmological data are sensitive




Normal Ordering

$$m_3 = \sqrt{\Delta m^2 + \delta m^2 / 2} = 5.06 \times 10^{-2} \text{ eV}$$




Oscillations independent on the absolute mass scale but give rise to a lower bound on Σ to which cosmological data are sensitive




Normal Ordering

$$m_3 = \sqrt{\Delta m^2 + \delta m^2/2} = 5.06 \times 10^{-2} \text{ eV}$$




$$\Sigma \gtrsim 6.5 \times 10^{-2} \text{ eV}$$

Oscillations independent on the absolute mass scale but give rise to a lower bound on Σ to which cosmological data are sensitive



Normal Ordering


$$m_3 = \sqrt{\Delta m^2 + \delta m^2/2} = 5.06 \times 10^{-2} \text{ eV}$$




$$\Sigma \gtrsim 6.5 \times 10^{-2} \text{ eV}$$

Inverted Ordering

$$m_2 = \sqrt{|\Delta m^2| + \delta m^2/2} = 5.04 \times 10^{-2} \text{ eV}$$




Oscillations independent on the absolute mass scale but give rise to a lower bound on Σ to which cosmological data are sensitive



Normal Ordering

$$m_3 = \sqrt{\Delta m^2 + \delta m^2/2} = 5.06 \times 10^{-2} \text{ eV}$$



$$\Sigma \gtrsim 6.5 \times 10^{-2} \text{ eV}$$

Inverted Ordering

$$m_2 = \sqrt{|\Delta m^2| + \delta m^2/2} = 5.04 \times 10^{-2} \text{ eV}$$




$$m_1 = \sqrt{|\Delta m^2| - \delta m^2/2} = 4.97 \times 10^{-2} \text{ eV}$$

$$m_3 = 0$$


$$\Sigma \gtrsim 10^{-1} \text{ eV}$$

Oscillations independent on the absolute mass scale but give rise to a lower bound on Σ to which cosmological data are sensitive



Normal Ordering

$$m_3 = \sqrt{\Delta m^2 + \delta m^2/2} = 5.06 \times 10^{-2} \text{ eV}$$



$$m_2 = \sqrt{\delta m^2} = 0.86 \times 10^{-2} \text{ eV}$$

$$m_1 = 0$$

$$\Sigma \gtrsim 6.5 \times 10^{-2} \text{ eV}$$

Inverted Ordering

$$m_2 = \sqrt{|\Delta m^2| + \delta m^2/2} = 5.04 \times 10^{-2} \text{ eV}$$





$$m_1 = \sqrt{|\Delta m^2| - \delta m^2/2} = 4.97 \times 10^{-2} \text{ eV}$$

$$m_3 = 0$$

$$\Sigma \gtrsim 10^{-1} \text{ eV}$$



The lower bound on Σ for IO only a factor ~2 smaller than the strongest limit set at present by cosmological data

Oscillations





Absolute minimum in NO, $\Delta\chi^2(\text{IO-NO}) = 3.6$

Oscillations





Absolute minimum in NO, $\Delta\chi^2(\text{IO-NO}) = 3.6$

Oscillations





Absolute minimum in NO, $\Delta\chi^2(\text{IO-NO}) = 3.6$

Oscillations



Absolute minimum in NO, $\Delta\chi^2(\text{IO-NO}) = 3.6$

Oscillations





Lower bounds on Σ

When minimising also with respect to the mass ordering
the allowed parameter space is the union of the contours

Absolute minimum in NO, $\Delta\chi^2(\text{IO-NO}) = 3.6$

Oscillations



Lower bounds on Σ

When minimising also with respect to the mass ordering
the allowed parameter space is the union of the contours

Absolute minimum in NO, $\Delta\chi^2(\text{IO-NO}) = 3.6$

Including $0\nu\beta\beta$ data


Including $0\nu\beta\beta$ data

KamLAND-Zen ^{136}Xe Limits (90% C.L.)

Phase 1 $T_{1/2}(0\nu) > 1.9 \times 10^{25} \text{ yr}$

Phase 2 $T_{1/2}(0\nu) > 9.2 \times 10^{25} \text{ yr}$

Combined $\mathbf{T_{1/2}(0\nu) > 1.07 \times 10^{26} \text{ yr}}$




Including $0\nu\beta\beta$ data

KamLAND-Zen ^{136}Xe Limits (90% C.L.)


Phase 1 $T_{1/2}(0\nu) > 1.9 \times 10^{25} \text{ yr}$

Phase 2 $T_{1/2}(0\nu) > 9.2 \times 10^{25} \text{ yr}$

Combined $\mathbf{T_{1/2}(0\nu) > 1.07 \times 10^{26} \text{ yr}}$





+NME Likelihood based on:
E.Lisi, A.Rotunno, F.Simkovic,
[arXiv:1506.04058](https://arxiv.org/abs/1506.04058)





$m_{\beta\beta} \lesssim 0.2 \text{ eV at } 2\sigma$



Oscill. + $0\nu\beta\beta$





Oscill. + $0\nu\beta\beta$



Oscill. + $0\nu\beta\beta$




Oscill. + $0\nu\beta\beta$



On the left: 2σ bound from KL-Zen when $m_{\beta\beta} \lesssim 0.2$ eV

Oscill. + $0\nu\beta\beta$



On the left: 2σ bound from KL-Zen when $m_{\beta\beta} \lesssim 0.2$ eV

On the right: constraint from KL-Zen added to the $\Delta\chi^2=3.6$ offset from oscillations \rightarrow stronger bound on $m_{\beta\beta}$ for IO

Cosmological Data

Cosmological Data

Two classes of models

Cosmological Data

Two classes of models

Λ CDM+ Σ : flat universe with a cosmological constant and adiabatic perturbations plus massive neutrinos (6+1 parameters)

Cosmological Data

Two classes of models

Λ CDM+ Σ : flat universe with a cosmological constant and adiabatic perturbations plus massive neutrinos (6+1 parameters)

Λ CDM+ Σ +Alens: one parameter more that controls lensing in the Planck angular spectra (6+1+1 parameters)

Cosmological Data

Two classes of models

Λ CDM+ Σ : flat universe with a cosmological constant and adiabatic perturbations plus massive neutrinos (6+1 parameters)

Λ CDM+ Σ +Alens: one parameter more that controls lensing in the Planck angular spectra (6+1+1 parameters)

We separately study NO and IO, taking as unknown the lowest neutrino mass and calculating the other two by means of the best-fit values of the mass square differences δm^2 and Δm^2

Cosmological Data

Two classes of models

Λ CDM+ Σ : flat universe with a cosmological constant and adiabatic perturbations plus massive neutrinos (6+1 parameters)

Λ CDM+ Σ +Alens: one parameter more that controls lensing in the Planck angular spectra (6+1+1 parameters)

We separately study NO and IO, taking as unknown the lowest neutrino mass and calculating the other two by means of the best-fit values of the mass square differences δm^2 and Δm^2

All in all $12 = 6 \times 2$ data set combinations

6 cases with $Alens=1$ and 6 with $Alens$ free

Planck TT + τ_{HFI}

Planck TT + τ_{HFI} + lensing

Planck TT + τ_{HFI} + BAO

Planck TT, TE, EE + τ_{HFI}

Planck TT, TE, EE + τ_{HFI} + lensing

Planck TT, TE, EE + τ_{HFI} + BAO

TT

TE,EE

τ_{HFI}

BAO

Temperature anisotropy

Polarization

Reionization prior on optical depth

Baryon acoustic oscillation

#	Model	Cosmological data set	$\Sigma/\text{eV (2}\sigma\text{)}, \text{NO}$	$\Sigma/\text{eV (2}\sigma\text{)}, \text{IO}$	$\Delta\chi^2_{\text{IO-NO}}$
1	$\Lambda\text{CDM} + \Sigma$	Planck TT + τ_{HFI}	< 0.72	< 0.80	0.7
2	$\Lambda\text{CDM} + \Sigma$	Planck TT + τ_{HFI} + lensing	< 0.64	< 0.63	0.2
3	$\Lambda\text{CDM} + \Sigma$	Planck TT + τ_{HFI} + BAO	< 0.21	< 0.23	1.2
4	$\Lambda\text{CDM} + \Sigma$	Planck TT, TE, EE + τ_{HFI}	< 0.44	< 0.48	0.6
5	$\Lambda\text{CDM} + \Sigma$	Planck TT, TE, EE + τ_{HFI} + lensing	< 0.45	< 0.47	0.3
6	$\Lambda\text{CDM} + \Sigma$	Planck TT, TE, EE + τ_{HFI} + BAO	< 0.18	< 0.20	1.6
7	$\Lambda\text{CDM} + \Sigma + A_{\text{lens}}$	Planck TT + τ_{HFI}	< 1.08	< 1.08	-0.1
8	$\Lambda\text{CDM} + \Sigma + A_{\text{lens}}$	Planck TT + τ_{HFI} + lensing	< 0.91	< 0.93	0.0
9	$\Lambda\text{CDM} + \Sigma + A_{\text{lens}}$	Planck TT + τ_{HFI} + BAO	< 0.45	< 0.46	0.2
10	$\Lambda\text{CDM} + \Sigma + A_{\text{lens}}$	Planck TT, TE, EE + τ_{HFI}	< 1.04	< 1.03	0.0
11	$\Lambda\text{CDM} + \Sigma + A_{\text{lens}}$	Planck TT, TE, EE + τ_{HFI} + lensing	< 0.89	< 0.89	0.1
12	$\Lambda\text{CDM} + \Sigma + A_{\text{lens}}$	Planck TT, TE, EE + τ_{HFI} + BAO	< 0.31	< 0.32	0.3

#	Model	Cosmological data set	$\Sigma/\text{eV (2}\sigma\text{)}, \text{NO}$	$\Sigma/\text{eV (2}\sigma\text{)}, \text{IO}$	$\Delta\chi^2_{\text{IO-NO}}$
1	$\Lambda\text{CDM} + \Sigma$	Planck TT + τ_{HFI}	< 0.72	< 0.80	0.7
2	$\Lambda\text{CDM} + \Sigma$	Planck TT + τ_{HFI} + lensing	< 0.64	< 0.63	0.2
3	$\Lambda\text{CDM} + \Sigma$	Planck TT + τ_{HFI} + BAO	< 0.21	< 0.23	1.2
4	$\Lambda\text{CDM} + \Sigma$	Planck TT, TE, EE + τ_{HFI}	< 0.44	< 0.48	0.6
5	$\Lambda\text{CDM} + \Sigma$	Planck TT, TE, EE + τ_{HFI} + lensing	< 0.45	< 0.47	0.3
6	$\Lambda\text{CDM} + \Sigma$	Planck TT, TE, EE + τ_{HFI} + BAO	< 0.18	< 0.20	1.6
7	$\Lambda\text{CDM} + \Sigma + A_{\text{lens}}$	Planck TT + τ_{HFI}	< 1.08	< 1.08	-0.1
8	$\Lambda\text{CDM} + \Sigma + A_{\text{lens}}$	Planck TT + τ_{HFI} + lensing	< 0.91	< 0.93	0.0
9	$\Lambda\text{CDM} + \Sigma + A_{\text{lens}}$	Planck TT + τ_{HFI} + BAO	< 0.45	< 0.46	0.2
10	$\Lambda\text{CDM} + \Sigma + A_{\text{lens}}$	Planck TT, TE, EE + τ_{HFI}	< 1.04	< 1.03	0.0
11	$\Lambda\text{CDM} + \Sigma + A_{\text{lens}}$	Planck TT, TE, EE + τ_{HFI} + lensing	< 0.89	< 0.89	0.1
12	$\Lambda\text{CDM} + \Sigma + A_{\text{lens}}$	Planck TT, TE, EE + τ_{HFI} + BAO	< 0.31	< 0.32	0.3

Some trends

#	Model	Cosmological data set	$\Sigma/\text{eV (2}\sigma\text{)}, \text{NO}$	$\Sigma/\text{eV (2}\sigma\text{)}, \text{IO}$	$\Delta\chi^2_{\text{IO-NO}}$
1	$\Lambda\text{CDM} + \Sigma$	Planck TT + τ_{HFI}	< 0.72	< 0.80	0.7
2	$\Lambda\text{CDM} + \Sigma$	Planck TT + τ_{HFI} + lensing	< 0.64	< 0.63	0.2
3	$\Lambda\text{CDM} + \Sigma$	Planck TT + τ_{HFI} + BAO	< 0.21	< 0.23	1.2
4	$\Lambda\text{CDM} + \Sigma$	Planck TT, TE, EE + τ_{HFI}	< 0.44	< 0.48	0.6
5	$\Lambda\text{CDM} + \Sigma$	Planck TT, TE, EE + τ_{HFI} + lensing	< 0.45	< 0.47	0.3
6	$\Lambda\text{CDM} + \Sigma$	Planck TT, TE, EE + τ_{HFI} + BAO	< 0.18	< 0.20	1.6
7	$\Lambda\text{CDM} + \Sigma + A_{\text{lens}}$	Planck TT + τ_{HFI}	< 1.08	< 1.08	-0.1
8	$\Lambda\text{CDM} + \Sigma + A_{\text{lens}}$	Planck TT + τ_{HFI} + lensing	< 0.91	< 0.93	0.0
9	$\Lambda\text{CDM} + \Sigma + A_{\text{lens}}$	Planck TT + τ_{HFI} + BAO	< 0.45	< 0.46	0.2
10	$\Lambda\text{CDM} + \Sigma + A_{\text{lens}}$	Planck TT, TE, EE + τ_{HFI}	< 1.04	< 1.03	0.0
11	$\Lambda\text{CDM} + \Sigma + A_{\text{lens}}$	Planck TT, TE, EE + τ_{HFI} + lensing	< 0.89	< 0.89	0.1
12	$\Lambda\text{CDM} + \Sigma + A_{\text{lens}}$	Planck TT, TE, EE + τ_{HFI} + BAO	< 0.31	< 0.32	0.3

Some trends

- Polarization and BAO data strengthen bounds on Σ

#	Model	Cosmological data set	$\Sigma/\text{eV (2}\sigma\text{)}, \text{NO}$	$\Sigma/\text{eV (2}\sigma\text{)}, \text{IO}$	$\Delta\chi^2_{\text{IO-NO}}$
1	$\Lambda\text{CDM} + \Sigma$	Planck TT + τ_{HFI}	< 0.72	< 0.80	0.7
2	$\Lambda\text{CDM} + \Sigma$	Planck TT + τ_{HFI} + lensing	< 0.64	< 0.63	0.2
3	$\Lambda\text{CDM} + \Sigma$	Planck TT + τ_{HFI} + BAO	< 0.21	< 0.23	1.2
4	$\Lambda\text{CDM} + \Sigma$	Planck TT, TE, EE + τ_{HFI}	< 0.44	< 0.48	0.6
5	$\Lambda\text{CDM} + \Sigma$	Planck TT, TE, EE + τ_{HFI} + lensing	< 0.45	< 0.47	0.3
6	$\Lambda\text{CDM} + \Sigma$	Planck TT, TE, EE + τ_{HFI} + BAO	< 0.18	< 0.20	1.6
7	$\Lambda\text{CDM} + \Sigma + A_{\text{lens}}$	Planck TT + τ_{HFI}	< 1.08	< 1.08	-0.1
8	$\Lambda\text{CDM} + \Sigma + A_{\text{lens}}$	Planck TT + τ_{HFI} + lensing	< 0.91	< 0.93	0.0
9	$\Lambda\text{CDM} + \Sigma + A_{\text{lens}}$	Planck TT + τ_{HFI} + BAO	< 0.45	< 0.46	0.2
10	$\Lambda\text{CDM} + \Sigma + A_{\text{lens}}$	Planck TT, TE, EE + τ_{HFI}	< 1.04	< 1.03	0.0
11	$\Lambda\text{CDM} + \Sigma + A_{\text{lens}}$	Planck TT, TE, EE + τ_{HFI} + lensing	< 0.89	< 0.89	0.1
12	$\Lambda\text{CDM} + \Sigma + A_{\text{lens}}$	Planck TT, TE, EE + τ_{HFI} + BAO	< 0.31	< 0.32	0.3

Some trends

- Polarization and BAO data strengthen bounds on Σ
- Bounds weaken up to a factor ~2 when A_{lens} free

#	Model	Cosmological data set	$\Sigma/\text{eV (}2\sigma\text{)}, \text{NO}$	$\Sigma/\text{eV (}2\sigma\text{)}, \text{IO}$	$\Delta\chi^2_{\text{IO-NO}}$
1	$\Lambda\text{CDM} + \Sigma$	Planck TT + τ_{HFI}	< 0.72	< 0.80	0.7
2	$\Lambda\text{CDM} + \Sigma$		< 0.64	< 0.63	0.2
3	$\Lambda\text{CDM} + \Sigma$		< 0.21	< 0.23	1.2
4	$\Lambda\text{CDM} + \Sigma$	Planck TT, TE, EE + τ_{HFI}	< 0.44	< 0.48	0.6
5	$\Lambda\text{CDM} + \Sigma$		< 0.45	< 0.47	0.3
6	$\Lambda\text{CDM} + \Sigma$		< 0.18	< 0.20	1.6
7	$\Lambda\text{CDM} + \Sigma + A_{\text{lens}}$	Planck TT + τ_{HFI}	< 1.08	< 1.08	-0.1
8	$\Lambda\text{CDM} + \Sigma + A_{\text{lens}}$	Planck TT + τ_{HFI} + lensing	< 0.91	< 0.93	0.0
9	$\Lambda\text{CDM} + \Sigma + A_{\text{lens}}$	Planck TT + τ_{HFI} + BAO	< 0.45	< 0.46	0.2
10	$\Lambda\text{CDM} + \Sigma + A_{\text{lens}}$	Planck TT, TE, EE + τ_{HFI}	< 1.04	< 1.03	0.0
11	$\Lambda\text{CDM} + \Sigma + A_{\text{lens}}$	Planck TT, TE, EE + τ_{HFI} + lensing	< 0.89	< 0.89	0.1
12	$\Lambda\text{CDM} + \Sigma + A_{\text{lens}}$	Planck TT, TE, EE + τ_{HFI} + BAO	< 0.31	< 0.32	0.3

Some trends

- Polarization and BAO data strengthen bounds on Σ
- Bounds weaken up to a factor ~2 when A_{lens} free
- Small differences between NO and IO
NO usually slightly preferred


#	Model	Cosmological data set	$\Sigma/\text{eV} (2\sigma)$, NO	$\Sigma/\text{eV} (2\sigma)$, IO	$\Delta\chi^2_{\text{IO-NO}}$
1	$\Lambda\text{CDM} + \Sigma$	Planck TT + τ_{HFI}	< 0.72	< 0.80	0.7
2	$\Lambda\text{CDM} + \Sigma$	Planck TT + τ_{HFI} + lensing	< 0.64	< 0.63	0.2
3	$\Lambda\text{CDM} + \Sigma$	Planck TT + τ_{HFI} + BAO	< 0.21	< 0.23	1.2
4	$\Lambda\text{CDM} + \Sigma$	Planck TT, TE, EE + τ_{HFI}	< 0.44	< 0.48	0.6
5	$\Lambda\text{CDM} + \Sigma$	Planck TT, TE, EE + τ_{HFI} + lensing	< 0.45	< 0.47	0.3
6	$\Lambda\text{CDM} + \Sigma$	Planck TT, TE, EE + τ_{HFI} + BAO	< 0.18	< 0.20	1.6
7	$\Lambda\text{CDM} + \Sigma + A_{\text{lens}}$	Planck TT + τ_{HFI}	< 1.08	< 1.08	-0.1
8	$\Lambda\text{CDM} + \Sigma + A_{\text{lens}}$	Planck TT + τ_{HFI} + lensing	< 0.91	< 0.93	0.0
9	$\Lambda\text{CDM} + \Sigma + A_{\text{lens}}$	Planck TT + τ_{HFI} + BAO	< 0.45	< 0.46	0.2
10	$\Lambda\text{CDM} + \Sigma + A_{\text{lens}}$	Planck TT, TE, EE + τ_{HFI}	< 1.04	< 1.03	0.0
11	$\Lambda\text{CDM} + \Sigma + A_{\text{lens}}$	Planck TT, TE, EE + τ_{HFI} + lensing	< 0.89	< 0.89	0.1
12	$\Lambda\text{CDM} + \Sigma + A_{\text{lens}}$	Planck TT, TE, EE + τ_{HFI} + BAO	< 0.31	< 0.32	0.3

Some trends

- Polarization and BAO data strengthen bounds on Σ
- Bounds weaken up to a factor ~2 when A_{lens} free
- Small differences between NO and IO
NO usually slightly preferred


Focus on 4 representative cases $\rightarrow (\#10, \#1, \#9, \#6)$

4 selected cases with increasingly strong bounds on Σ



4 selected cases with increasingly strong bounds on Σ


- χ^2 curves for NO and IO converge for large Σ



4 selected cases with increasingly strong bounds on Σ


- χ^2 curves for NO and IO converge for large Σ

- χ^2 curves bifurcate for small Σ




4 selected cases with increasingly strong bounds on Σ

- χ^2 curves for NO and IO converge for large Σ
- χ^2 curves bifurcate for small Σ
- $\Sigma = 0$ not allowed



4 selected cases with increasingly strong bounds on Σ



- χ^2 curves for NO and IO converge for large Σ
- χ^2 curves bifurcate for small Σ
- $\Sigma = 0$ not allowed
- For cases #10 and #9 the minimum of the χ^2 is reached for a value of Σ higher than the minimum allowed




Combination of oscillation and non-osculation data

Combination of oscillation and non-osculation data

Oscillations





Combination of oscillation and non-osculation data





Combination of oscillation and non-osculation data

Oscill. + $0\nu\beta\beta$ + Cosmo #10





Combination of oscillation and non-osculation data

Oscill. + $0\nu\beta\beta$ + Cosmo #1





Combination of oscillation and non-osculation data

Oscill. + $0\nu\beta\beta$ + Cosmo #9



Combination of oscillation and non-osculation data

Oscill. + $0\nu\beta\beta$ + Cosmo #6



Conclusions

Conclusions

- Ranges of well-known 3v parameters ($\delta m^2, \theta_{12}$) & ($\Delta m^2, \theta_{13}$) confirmed by v2017 data updates

Conclusions

- Ranges of well-known 3v parameters ($\delta m^2, \theta_{12}$) & ($\Delta m^2, \theta_{13}$) confirmed by v2017 data updates
- CPV: $\sin\delta < 0$ preferred

Conclusions

- Ranges of well-known 3v parameters ($\delta m^2, \theta_{12}$) & ($\Delta m^2, \theta_{13}$) confirmed by v2017 data updates
- CPV: $\sin\delta < 0$ preferred
 - best fit: $\delta/\pi \sim 1.3 - 1.4 \pm 0.2$ (1σ)
 - $\sin\delta \sim 0$ disfavoured at $> 2\sigma$
 - $\sin\delta \sim +1$ disfavoured at $> 3\sigma$

Conclusions

- Ranges of well-known 3v parameters ($\delta m^2, \theta_{12}$) & ($\Delta m^2, \theta_{13}$) confirmed by v2017 data updates
- CPV: $\sin\delta < 0$ preferred
 - best fit: $\delta/\pi \sim 1.3 - 1.4 \pm 0.2$ (1σ)
 - $\sin\delta \sim 0$ disfavoured at $> 2\sigma$
 - $\sin\delta \sim +1$ disfavoured at $> 3\sigma$
- Octant info: still fragile and dependent on mass ordering

Conclusions

- Ranges of well-known 3ν parameters ($\delta m^2, \theta_{12}$) & ($\Delta m^2, \theta_{13}$) confirmed by v2017 data updates
- CPV: $\sin\delta < 0$ preferred
 - best fit: $\delta/\pi \sim 1.3 - 1.4 \pm 0.2$ (1σ)
 - $\sin\delta \sim 0$ disfavoured at $> 2\sigma$
 - $\sin\delta \sim +1$ disfavoured at $> 3\sigma$
- Octant info: still fragile and dependent on mass ordering
- Mass Ordering: IO disfavored by oscillation data:

	LBL+SOL+KL	+SBL	+ATM
$\Delta\chi^2(\text{IO-NO})$	1.1	1.1	3.6

Conclusions

- Ranges of well-known 3ν parameters ($\delta m^2, \theta_{12}$) & ($\Delta m^2, \theta_{13}$) confirmed by v2017 data updates
- CPV: $\sin\delta < 0$ preferred
 - best fit: $\delta/\pi \sim 1.3 - 1.4 \pm 0.2$ (1σ)
 - $\sin\delta \sim 0$ disfavoured at $> 2\sigma$
 - $\sin\delta \sim +1$ disfavoured at $> 3\sigma$
- Octant info: still fragile and dependent on mass ordering
- Mass Ordering: IO disfavored by oscillation data:

	LBL+SOL+KL	+SBL	+ATM
$\Delta\chi^2(\text{IO-NO})$	1.1	1.1	3.6
- Non oscillation data corroborate NO ($\Delta\chi^2_{\text{IO-NO}} \in [3.6, 4.4]$)